



Can a Probiotic Help to Prevent Travelers Diarrhea?

By: Peter J. Whorwell, M.D., Ph.D., Director, South Manchester Functional Bowel Service, Wythenshawe Hospital and Professor, Institute of Inflammation and Repair, University of Manchester, Manchester, United Kingdom

QUESTION – I will soon be travelling to several countries where finding safe drinking water will be a challenge. In the event I eat or drink something that would cause traveler’s diarrhea would it be wise to preventatively start taking a probiotic? Is it alright to take loperamide at the same time as taking a probiotic?

ANSWER – There is some evidence that taking a probiotic can help prevent travellers’ diarrhea although probiotics are notoriously variable in terms of their activity, with some being more effective than others with respect to this indication. I see no reason why you shouldn’t take loperamide with a probiotic should the need arise.

I should point out that diarrhea in gastroenteritis does serve a purpose of clearing out the germs that are causing the problem and, therefore, taking loperamide can sometimes delay recovery.

I usually advise patients to take one of the glucose and electrolyte powders when they get travellers’ diarrhea and just use the loperamide for emergencies, travelling, and to get home.

Suggested IFFGD Reading

Thompson, W. *Managing Diarrhea*. IFFGD. Fact Sheet No. 201.

Bonci, L. *Nutrition Strategies for Managing Diarrhea*. IFFGD. Fact Sheet No. 208.

The information in this fact sheet was adapted from the Clinical Corner Section of Digestive Health Matters, Vol. 18, No. 2.

About IFFGD

The International Foundation for Functional Gastrointestinal Disorders (IFFGD) is a 501(c)(3) nonprofit education and research organization. We work to promote awareness, scientific advancement, and improved care for people affected by chronic digestive conditions. Our mission is to inform, assist, and support people affected by gastrointestinal disorders. Founded in 1991, we rely on donors to carry out our mission. Visit our websites at: www.iffgd.org or www.aboutIBS.org.

About the Publication

Opinions expressed are an author’s own and not necessarily those of the International Foundation for Functional Gastrointestinal Disorders (IFFGD). IFFGD does not guarantee or endorse any product in this publication or any claim made by an author and disclaims all liability relating thereto. This article is in no way intended to replace the knowledge or diagnosis of your doctor. We advise seeing a physician whenever a health problem arises requiring an expert’s care.

For more information, or permission to reprint this article, contact IFFGD by phone at 414-964-1799 or by email at iffgd@iffgd.org.