



GERD, Hiatal Hernia, and Surgery

By: J. Patrick Waring, M.D., Digestive Healthcare of Georgia, Atlanta, Georgia

523



International Foundation for Gastrointestinal Disorders (www.iffgd.org)



Reading time: 8 minutes

© Copyright 2002-2006 by the International Foundation for Gastrointestinal Disorders

Question:

I have GERD and have been told that I may need surgery to repair a hiatal hernia. Can you please explain the surgery? Will my GERD be resolved? What are the potential risks related to the surgery?

Answer:

Since its introduction around 10 years ago, laparoscopic surgery for gastroesophageal reflux disease (GERD) has become one of the more common operations performed in the United States. It is interesting that there is still not a consensus amongst physicians as to the proper role of this procedure in the management of GERD. Many physicians (especially surgeons) are very fond of this operation. However, many physicians (mostly non-surgeons) are very much opposed to it. Consequently, patients are often faced with information from different sources that appears to be completely contradictory.

The basic problem in patients with GERD is that acid from the stomach refluxes up into the esophagus (food pipe). The barrier that prevents this from happening is complex. However, there are only a couple of things that can go wrong here. First, the muscle at the bottom of the esophagus – the lower esophageal sphincter – may become weakened. Second, one may have a hiatal hernia. (In health, the esophagus should be in the chest and the stomach should be in the abdomen. The diaphragm is a thin muscle that separates the chest from the abdomen. In patients with a hiatal hernia, the stomach has moved up above the diaphragm and into the chest.) When either of these problems happens, the barrier between the stomach and the esophagus is less competent and more likely to allow reflux of stomach acid into the esophagus. A hiatal hernia tends to make the anti-reflux barrier more open and weaker. Consequently, it allows acid reflux to occur more easily. The hiatal hernia itself usually does not cause symptoms unless it is very large.

Most of the data on success rates comes from referral centers where this type of surgery is performed every day. At USC Medical Center in Los Angeles, one hundred patients were followed for an average of 21 months. Ninety-six percent of patients had a good outcome, and 71% were completely free of symptoms. Only 3 patients were back on medical therapy; 2 patients were worsened by the procedure. The same results are not seen in less experienced centers. A recent study

showed only a 57% satisfaction rate, with 67% of the patients developing new symptoms, such as difficulty swallowing or bloating. Additionally, 6.7% of the patients needed a repeat operation in less than one year.

Laparoscopic Surgery

The surgical procedures to treat GERD may be performed through the abdomen or the chest and can be done open or laparoscopically. Laparoscopic surgery is minimally invasive and has certain postoperative advantages. It is important to be aware that “minimally invasive” describes the surgical technique. It is called minimally invasive because of the small incisions that can be used. A laparoscopic fundoplication is major surgery and the patient and surgeon should think of it in that light. Only an experienced surgeon who has done many laparoscopic operations should perform it. Laparoscopic surgery has many of the risks for major complications that open surgery entails. The primary advantage of minimally invasive surgery involves the degree of patient discomfort following the surgery and the patient’s recovery time; both are decreased following laparoscopic surgery when compared to open surgery.

Surgical therapy corrects these underlying problems. The surgeon can pull the stomach back into the abdomen, correcting the hiatal hernia. Furthermore, whether or not a hiatal hernia is present, the surgeon can take the top part of the stomach (the fundus) and wrap it around the lower esophagus, much like a tailor can cuff a pair of pants (a procedure called fundoplication). This strengthens the lower esophageal sphincter muscle and helps prevent acid reflux from occurring. The procedure can nearly always be done laparoscopically, using very small incisions. Typically, patients will spend the next one to two days in the hospital and 2–4 weeks at home recovering. Ideally, within a few weeks, patients will no longer have heartburn, and no longer need to take heartburn medication. The most common complications are difficulty swallowing, abdominal bloating, diarrhea, and nausea. Most patients can’t belch as well as they could before surgery, although the inability to belch is distinctly uncommon. About 25% of patients can’t vomit after surgery.

Longer follow-up data is available from the Mayo Clinic in Jacksonville, Florida. Again, 96% were satisfied with their results between 5 and 8 years later. Only 14% were back on medical therapy, 6% had symptoms of chest pain, 6% had

heartburn and 6% had regurgitation (the feeling of fluid coming back up into the esophagus). Twenty percent of the patients had abdominal bloating, 12% had diarrhea, 27% reported some difficulty swallowing, and 7% required a dilation procedure to stretch the esophagus. Again, the same results may not apply to less experienced centers. A recently published study from the Veterans Administration showed that 62% of patients having surgery required medications for heartburn after 10–13 years.

There are some special situations that deserve mention. It appears that the ideal candidate for the operation is the patient with heartburn that responds nicely to medical therapy. Patients with atypical symptoms of GERD, such as laryngitis, chronic cough, or asthma don't seem to do quite as well. Patients with a poor response to medical therapy may not do as well either. The main reason for this is that these patients may have problems besides GERD that are causing these *refractory* symptoms [symptoms that are unresponsive to treatment]. Some patients develop a stricture, which is a narrowing of the esophagus caused by severe acid reflux. The surgical results in these patients are not quite as good as in patients who don't have a stricture. Having surgery for a stricture is still controversial, but it is my belief that most patients with this problem are better off not having surgery.

Another controversial area is Barrett's esophagus. This is a situation where the lining of the esophagus changes to a potentially pre-malignant tissue. Some surgeons feel strongly that surgery will decrease the risk of developing cancer, and will so inform their patients. Unfortunately, there is no compelling scientific evidence that this is true. Most experts in this area believe that surgery should not be performed for the sole purpose of preventing esophageal cancer.

There are some patients with a very large hiatal hernia, where one-half of the stomach or more is actually in the chest. The currently available medical therapy rarely works in these patients and surgery is usually necessary.

Surgery for GERD should be considered in three circumstances. First, patients who have side effects to medical therapy may require surgery. However, please keep in mind that there are now five proton pump inhibitors. It is uncommon to see a patient who cannot tolerate any of the five. Second, patients who are poorly responsive to medical therapy may benefit from surgery. However, as stated above, some refractory symptoms may be unrelated to GERD. The physician should be as certain as possible that the symptoms are indeed due to GERD. Third, patients who wish to be free of the need to take medications for their GERD may choose to have surgery. [Note that some patients must continue to use anti-reflux medication even after surgery.]

Patients must be aware that the overall satisfaction rate is around 95%, and the chance of being symptom-free is about 70%. I rarely have a patient ask me about what happens to patients who don't do well; they seem to assume that they will

be one of the 95% who have a good outcome. Many of these patients simply don't respond and need to continue taking medication. However, 1–2% of all surgical patients are worse after the procedure. It must be pointed out that the few patients who do poorly with this operation have a life-altering experience. They often have difficulty with swallowing, nausea, chest pain, or abdominal pain that is refractory to all medical and surgical therapies and occurs with every meal for the rest of their lives.

Over ten years of experience has taught us several important lessons. First, there is no substitute for surgical experience. If you are considering surgery for the treatment of GERD, ask your doctor how many of these operations they have done. The best results can be expected from surgeons who have done 200–400 operations or more. Second, the patients who do the best are those with simple heartburn that responds nicely to medical therapy. If your symptoms don't respond to medical treatment, ask your doctor about how certain he or she is that the symptoms are related to GERD and not something else. Third, several diagnostic tests are imperative before surgery. Make sure to ask your doctor if all necessary testing is completed and that the results suggest that a good outcome will be obtained with a successful operation. If these criteria are met, the chances are favorable for a good, long-term response to surgical therapy for GERD.

This information is provided for general information and is not intended to replace your doctor's advice.

About IFFGD

The International Foundation for Gastrointestinal Disorders (IFFGD) is a 501(c)(3) nonprofit education and research organization. We work to promote awareness, scientific advancement, and improved care for people affected by chronic digestive conditions. Our mission is to inform, assist, and support people affected by gastrointestinal disorders. Founded in 1991, we rely on donors to carry out our mission. Visit our website at: www.iffgd.org or www.aboutgerd.org.

IFFGD

537 Long Point Road, Suite 101
Mt Pleasant, SC 29464

About the Publication

Opinions expressed are an author's own and not necessarily those of the International Foundation for Gastrointestinal Disorders (IFFGD). IFFGD does not guarantee or endorse any product in this publication or any claim made by an author and disclaims all liability relating thereto. This article is in no way intended to replace the knowledge or diagnosis of your healthcare provider. We advise seeing a healthcare provider whenever a health problem arises requiring an expert's care.

For more information, or permission to reprint this article, contact IFFGD by phone at 414-964-1799 or by email at iffgd@iffgd.org.