



Gastric Antral Vascular Ectasia (Watermelon Stomach)

570



By: Shajan Peter, MD, Division of Gastroenterology and Hepatology,
Department of Medicine, University of Alabama at Birmingham, AL

International Foundation for Gastrointestinal Disorders (www.iffgd.org)

© Copyright 2019 by the International Foundation for Gastrointestinal Disorders

Understanding this Syndrome and Treatment Options

What is GAVE?

Gastric Antral Vascular Ectasia (GAVE), sometimes called “watermelon stomach,” is a rare condition in which the blood vessels of the stomach lining become fragile and leak blood. When seen through an endoscope (an instrument physicians use to view the inside of the esophagus and stomach), a GAVE stomach resembles the markings on watermelons. GAVE may cause chronic anemia and sometimes other signs of blood loss such as blood in the stool.

What causes GAVE?

The specific cause of GAVE is unknown. However, it is often associated with cirrhosis of the liver. It can also be linked with other conditions such as autoimmune disorders, chronic renal failure, cardiac disease, diabetes mellitus, atrophic gastritis, cirrhosis, scleroderma, pernicious anemia and CREST syndrome. It can also occur in patients with portal hypertension, vascular disease, metabolic syndrome.

What are symptoms of GAVE?

GAVE is characterized primarily by gastrointestinal bleeding, which may result in

- fatigue caused by anemia (a condition that develops when your blood lacks enough healthy red blood cells)
- Hematemesis (vomiting blood)
- Blood in the stools

How is GAVE diagnosed?

Endoscopic examination is the primary diagnostic test. Seen through the endoscope, GAVE stomachs have stripes of redness (“watermelon appearance”), or show honeycombing or nodules. Frequently, those areas ooze blood during the endoscopic examination. Tissue biopsy can aid in the diagnosis and can show characteristic findings such as dilated blood vessels among other features.

How is GAVE treated?

GAVE may cause no symptoms; however, it can cause chronic blood loss anemia, requiring iron replacement or frequent blood transfusions. Various endoscopic treatments can be used and, currently, are the best way to prevent further blood loss. These treatments cause scarring of the abnormal areas which reduces the oozing and bleeding. This will decrease the need for blood transfusions and will stabilize the blood count.

Besides endoscopic treatment, hormonal and other drug therapies have been tried with varying success. A very few patients may need surgery. Treatment of the underlying conditions associated with GAVE, such as liver disease and diabetes can help minimize GAVE. In patients with cirrhosis, liver transplantation may eliminate GAVE.

About IFFGD

The International Foundation for Gastrointestinal Disorders (IFFGD) is a 501(c)(3) nonprofit education and research organization. We work to promote awareness, scientific advancement, and improved care for people affected by chronic digestive conditions. Our mission is to inform, assist, and support people affected by gastrointestinal disorders. Founded in 1991, we rely on donors to carry out our mission. Visit our website at: www.iffgd.org.

IFFGD

537 Long Point Road, Suite 101
Mt Pleasant, SC 29464

About the Publication

Opinions expressed are the author’s own and not necessarily those of the International Foundation for Gastrointestinal Disorders (IFFGD). IFFGD does not guarantee or endorse any product in this publication or any claim made by the author and disclaims all liability relating thereto. This article is in no way intended to replace the knowledge or diagnosis of your doctor. We advise seeing a physician whenever a health problem arises requiring an expert’s care.

For more information, or permission to reprint this article, contact IFFGD by phone at 414-964-1799 or by email at iffgd@iffgd.org

