



Report from IFFGD Research Award Winner:
**Biopsychosocial Perspectives on Assessment
and Treatment**

By: Carolina Elder Danda, Ph.D, LLC., Prairie Village, KS

831



International Foundation for Gastrointestinal Disorders (www.iffgd.org)

 Reading time: 4 minutes © Copyright 2003-2007 by the International Foundation for Gastrointestinal Disorders

Dr. Danda is the recipient of the **IFFGD 2003 Research Award to Pediatric Investigator, Clinical Science**. She received her Bachelor's degree in Psychology and French from the University of Missouri-Columbia in 1993 and attended graduate school at the University of Florida Health Sciences Center in the Department of Clinical and Health Psychology, where she specialized in pediatric psychology, participated in numerous independent and collaborative research projects, and had several publications. She continued training as an intern at Children's Mercy Hospital and a postdoctoral fellow at the University of Kansas Medical Center in the Department of Pediatrics before establishing her private practice in Prairie Village, Kansas.

As many as 10–20% of school-age children experience bellyaches severe enough to affect activities. Not all of these children seek medical treatment or experience chronic, continuous abdominal pain. A subset of children present to their clinician with symptoms sufficient to meet criteria for a functional gastrointestinal disorder (e.g., functional dyspepsia, irritable bowel syndrome). Past research on childhood abdominal pain used a broad description – *recurrent abdominal pain (RAP)*. Rome II criteria, however, classifies abdominal pain into subsets based on symptom presentation. My research focuses on examining the psychosocial contributions to these functional gastrointestinal disorders and determining whether there are different psychosocial factors (e.g., emotional or behavioral symptoms, developmental stressors, family communication patterns) associated with the specific functional gastrointestinal disorders.

I am particularly interested in identifying factors that predispose children and adolescents with functional gastrointestinal disorders to experience disability and identifying psychosocial factors that affect treatment outcome. I am collaborating on a research project

collecting data to provide a description of the developmental, psychosocial, and social factors influencing children with functional abdominal pain disorders and compare medical (e.g., drugs such as imipramine, montelukast, ranitidine, and erythromycin) and psychological (e.g., cognitive behavioral therapy) treatment for functional dyspepsia. In addition to comparing treatment modalities, we hope to identify factors associated with treatment outcome. Essentially, we want to find out which treatments work for which patients. Treatment of functional gastrointestinal disorders does not appear to be “one size fits all.”

Another core issue related to functional gastrointestinal disorders is disability. Some children experience interruption in their daily activities due to abdominal pain, whereas others do not. Which children experience disability related to a functional gastrointestinal disorder and what factors predispose them to experience disability?

Pain Associated Disability Syndrome (PADS) is a term used to describe patients who experience a downward spiral of disability (i.e., school absence, withdrawal from friends, restriction of daily activities) and pain for which acute pain management strategies have failed. A retrospective analysis [examination of known outcomes using existing records] indicated that almost all patients with PADS had a functional gastrointestinal disorder as well as physical and emotional stressors that sustained pain and arousal. Pain and disability are exacerbated or maintained via medical, psychosocial, and developmental factors. For example, pain experiences can lead to an increased sensitivity to potentially painful stimuli (visceral hyperalgesia). Undergoing multiple medical tests without explanation can lead to heightened anxiety and arousal. Developmental challenges, such as transitions to a new school, disruptions in a peer group,

and puberty, can lead to heightened states of stress. Heightened states of stress and arousal, in turn, can exacerbate pain sensation and focus.

In collaboration with other team members here at the University of Kansas, I plan to systematically and prospectively define the characteristics of patients with PADS, examine the effects of multidisciplinary treatment, and identify factors associated with treatment outcome. One area I plan to explore is the contribution of anxiety related to somatic symptoms and generalized anxiety or depression. Parents and children's negative or worrisome thoughts about medical problems are likely to play a large role in the development of PADS. For example, many children with a functional gastrointestinal disorder undergo multiple medical tests that result in negative results, which then increase anxiety, either in the patient and/or the parents. In addition, an anxiety or depressive disorder may lead to a predisposition to experience somatic symptoms, such as abdominal pain. Family dynamics and communication, especially during adolescent years, can also play a role in exacerbating and maintaining pain and disability. Research on PADS patients will ultimately help provide information to diagnose and treat these youths more effectively.

Lastly, I am involved in several projects pertaining to *Functional Fecal Retention (FFR)*, a common diagnosis associated with constipation that is characterized by at least 12 weeks of passage of large diameter stools at intervals less than two times a week; and retentive posturing (i.e., avoiding defecation by purposefully contracting the pelvic floor). Presumably, children with constipation experience pain with defecation of large, hard stools and develop a fear of defecation. Toilet learning is either failed or interrupted, and children appear unwilling or unable to relax the pelvic floor, creating an obstruction to defecation. At the University of Kansas, we are examining whether fear actually increases anal sphincter pressure and whether imagery and relaxation reduce fear and anal sphincter pressure. Children are randomized to either no treatment or treatment involving learning relaxation and imagery. Physiological and visual analog readings of anxiety are taken before, during, and after colonic manometry catheter removal.

Another study involves piloting a developmentally friendly, educational, non-invasive treatment of FFR. FFR may often go unrecognized at first, and children may be subjected to invasive and painful procedures, including enemas and repeated rectal exams, that could increase fear of defecation. The new approach to treatment includes

education for the child and family about FFR, stool disimpaction with MiraLax, and assurance of painless defecation, as well as brief behavioral recommendations about differential attention to toileting behaviors (i.e., praising appropriate toileting and other positive behaviors while minimizing attention to problematic behaviors). Goals of the study include determining the effect of this first, primarily educational visit on symptom resolution and identifying medical and psychosocial factors associated with treatment success and failure. Children with encopresis, or fecal soiling, tend to have more behavioral and social problems than other children, although not necessarily at clinical levels. The question remains, however, what role does oppositional behavior, attention problems, anxiety, or depressive symptoms play in children with persistent FFR? If children have coexisting psychological problems, does treatment of these disorders lead to improvements in FFR?

About IFFGD

The International Foundation for Functional Gastrointestinal Disorders (IFFGD) is a 501(c)(3) nonprofit education and research organization. We work to promote awareness, scientific advancement, and improved care for people affected by chronic digestive conditions. Our mission is to inform, assist, and support people affected by gastrointestinal disorders. Founded in 1991, we rely on donors to carry out our mission. Visit our websites at: www.iffgd.org or www.aboutKidsGI.org.

IFFGD

537 Long Point Rd., Suite 101
Mt. Pleasant, S.C, 29464

About the Publication

Opinions expressed are an author's own and not necessarily those of the International Foundation for Functional Gastrointestinal Disorders (IFFGD). IFFGD does not guarantee or endorse any product in this publication or any claim made by an author and disclaims all liability relating thereto. This article is in no way intended to replace the knowledge or diagnosis of your doctor. We advise seeing a physician whenever a health problem arises requiring an expert's care.

For more information, or permission to reprint this article, contact IFFGD by phone at 414-964-1799 or by email at iffgd@iffgd.org.
